

417

418

Table 3

419

Incidence of Adverse Events \geq 5% of Patients in Clinical Trials (N =

420

356)

421

(Adverse Events were followed for a period of 12 months following

422

RITUXAN therapy.)

	All Grades (%)	Grade 3 and 4 (%)
Any Adverse Events	99	57
Body as a Whole	86	10
Fever	53	1
Chills	33	3
Infection	31	4
Asthenia	26	1
Headache	19	1
Abdominal Pain	14	1
Pain	12	1
Back Pain	10	1
Throat Irritation	9	0
Flushing	5	0
Cardiovascular System	25	3
Hypotension	10	1
Hypertension	6	1
Digestive System	37	2
Nausea	23	1
Diarrhea	10	1
Vomiting	10	1
Hemic and Lymphatic System	67	48
Lymphopenia	48	40
Leukopenia	14	4
Neutropenia	14	6
Thrombocytopenia	12	2
Anemia	8	3
Metabolic and Nutritional Disorders	38	3
Angioedema	11	1
Hyperglycemia	9	1
Peripheral Edema	8	0
LDH Increase	7	0
Musculoskeletal System	26	3
Myalgia	10	1
Arthralgia	10	1
Nervous System	32	1
Dizziness	10	1
Anxiety	5	1
Respiratory System	38	4
Increased Cough	13	1
Rhinitis	12	1

	All Grades (%)	Grade 3 and 4 (%)
Bronchospasm	8	1
Dyspnea	7	1
Sinusitis	6	0
Skin and Appendages	44	2
Night Sweats	15	1
Rash	15	1
Pruritus	14	1
Urticaria	8	1

423
424
425

426 **Risk Factors Associated with Increased Rates of Adverse Events:**

427 Administration of RITUXAN weekly for 8 doses resulted in higher rates of
428 Grade 3 and 4 adverse events¹⁷ overall (70%) compared with
429 administration weekly for 4 doses (57%). The incidence of Grade 3 or 4
430 adverse events was similar in patients retreated with RITUXAN compared
431 with initial treatment (58% and 57%, respectively). The incidence of the
432 following clinically significant adverse events was higher in patients with
433 bulky disease (lesions ≥ 10 cm) (N = 39) versus patients with lesions
434 < 10 cm (N = 195): abdominal pain, anemia, dyspnea, hypotension, and
435 neutropenia.

436

437 **Infusion Reactions (See BOXED WARNINGS and WARNINGS):** Mild to
438 moderate infusion reactions consisting of fever and chills/rigors occurred
439 in the majority of patients during the first RITUXAN infusion. Other
440 frequent infusion reaction symptoms included nausea, pruritus,
441 angioedema, asthenia, hypotension, headache, bronchospasm, throat
442 irritation, rhinitis, urticaria, rash, vomiting, myalgia, dizziness, and

443 hypertension. These reactions generally occurred within 30 to 120
444 minutes of beginning the first infusion, and resolved with slowing or
445 interruption of the RITUXAN infusion and with supportive care
446 (diphenhydramine, acetaminophen, IV saline, and vasopressors). In an
447 analysis of data from 356 patients with relapsed or refractory, low-grade
448 NHL who received 4 (N = 319) or 8 (N = 37) weekly infusions of
449 RITUXAN, the incidence of infusion reactions was highest during the first
450 infusion (77%) and decreased with each subsequent infusion (30% with
451 fourth infusion and 14% with eighth infusion).

452

453 **Infectious Events:** RITUXAN induced B-cell depletion in 70% to 80% of
454 patients and was associated with decreased serum immunoglobulins in a
455 minority of patients; the lymphopenia lasted a median of 14 days (range, 1
456 to 588 days). Infectious events occurred in 31% of patients: 19% of
457 patients had bacterial infections, 10% had viral infections, 1% had fungal
458 infections, and 6% were unknown infections. Incidence is not additive
459 because a single patient may have had more than one type of infection.
460 Serious infectious events (Grade 3 or 4),¹⁷ including sepsis, occurred in
461 2% of patients.

462

463 **Hematologic Events:** Grade 3 and 4 cytopenias¹⁷ were reported in 12%
464 of patients treated with RITUXAN; these include: lymphopenia (40%),
465 neutropenia (6%), leukopenia (4%), anemia (3%), and thrombocytopenia

(2%). The median duration of lymphopenia was 14 days (range, 1 to 588 days) and of neutropenia was 13 days (range, 2 to 116 days). A single occurrence of transient aplastic anemia (pure red cell aplasia) and two occurrences of hemolytic anemia following RITUXAN therapy were reported. In addition, there have been rare postmarketing reports of prolonged pancytopenia and marrow hypoplasia.

Cardiac Events (See BOXED WARNINGS): Grade 3 or 4 cardiac-related events include hypotension. Rare, fatal cardiac failure with symptomatic onset weeks after RITUXAN has also been reported. Patients who develop clinically significant cardiopulmonary events should have RITUXAN infusion discontinued.

Pulmonary Events (See BOXED WARNINGS): 135 patients (38%) experienced pulmonary events. The most common respiratory system adverse events experienced were increased cough, rhinitis, bronchospasm, dyspnea, and sinusitis. Three pulmonary events have been reported in temporal association with RITUXAN infusion as a single agent: acute bronchospasm, acute pneumonitis presenting 1–4 weeks post-RITUXAN infusion, and bronchiolitis obliterans. One case of bronchiolitis obliterans was associated with progressive pulmonary symptoms and culminated in death several months following the last RITUXAN infusion. The safety of resumption or continued administration

489 of RITUXAN in patients with pneumonitis or bronchiolitis obliterans is
490 unknown.

491

492 **Immune/Autoimmune Events:** Immune/autoimmune events have been
493 reported, including uveitis, optic neuritis in a patient with systemic
494 vasculitis, pleuritis in a patient with a lupus-like syndrome, serum sickness
495 with polyarticular arthritis, and vasculitis with rash.

496

497 **Less Commonly Observed Events:** In clinical trials, < 5% and > 1% of
498 the patients experienced the following events regardless of causality
499 assessment:

500 agitation, anorexia, arthritis, conjunctivitis, depression, dyspepsia, edema,
501 hyperkinesia, hypertonia, hypesthesia, hypoglycemia, injection site pain,
502 insomnia, lacrimation disorder, malaise, nervousness, neuritis,
503 neuropathy, paresthesia, somnolence, vertigo, weight decrease.

504

505 **OVERDOSAGE**

506 There has been no experience with overdosage in human clinical trials.

507 Single doses of up to 500 mg/m² have been given in controlled clinical
508 trials.¹⁰

509

510 **DOSAGE AND ADMINISTRATION**

511 **Initial Therapy:**

512 RITUXAN is given at 375 mg/m² IV infusion once weekly for 4 or 8 doses.

513

514 **Retreatment Therapy:**

515 Patients who subsequently develop progressive disease may be safely

516 retreated with RITUXAN 375 mg/m² IV infusion once weekly for 4 doses.

517 Currently there are limited data concerning more than 2 courses.

518

519 **RITUXAN as a Component of Zevalin (Ibritumomab Tiuxetan)**

520 **Therapeutic Regimen:**

521 As a required component of the Zevalin therapeutic regimen, RITUXAN

522 250 mg/m² should be infused within 4 hours prior to the administration of

523 Indium-111- (In-111-) Zevalin and within 4 hours prior to the

524 administration of Yttrium-90- (Y-90-) Zevalin. Administration of RITUXAN

525 and In-111-Zevalin should precede RITUXAN and Y-90-Zevalin by 7-9

526 days. Refer to the Zevalin package insert for full prescribing information

527 regarding the Zevalin therapeutic regimen.

528

529 RITUXAN may be administered in an outpatient setting. **DO NOT**

530 **ADMINISTER AS AN INTRAVENOUS PUSH OR BOLUS. (See**

531 **Administration.)**

532

533 **Instructions for Administration**

534 **Preparation for Administration:** Use appropriate aseptic technique.

535 Withdraw the necessary amount of RITUXAN and dilute to a final

536 concentration of 1 to 4 mg/mL into an infusion bag containing either

537 0.9% Sodium Chloride, USP, or 5% Dextrose in Water, USP. Gently

538 invert the bag to mix the solution. Discard any unused portion left in the

539 vial. Parenteral drug products should be inspected visually for particulate

540 matter and discoloration prior to administration.

541

542 RITUXAN solutions for infusion may be stored at 2–8°C (36–46°F) for 24

543 hours. RITUXAN solutions for infusion have been shown to be stable for

544 an additional 24 hours at room temperature. However, since RITUXAN

545 solutions do not contain a preservative, diluted solutions should be stored

546 refrigerated (2–8°C). No incompatibilities between RITUXAN and

547 polyvinylchloride or polyethylene bags have been observed.

548

549 **Administration: DO NOT ADMINISTER AS AN INTRAVENOUS PUSH**

550 **OR BOLUS.** Infusion and hypersensitivity reactions may occur (see

551 **BOXED WARNINGS, WARNINGS, and ADVERSE REACTIONS).**

552 Premedication consisting of acetaminophen and diphenhydramine should

553 be considered before each infusion of RITUXAN. Premedication may

554 attenuate infusion reactions. Since transient hypotension may occur

555 during RITUXAN infusion, consideration should be given to withholding

556 antihypertensive medications 12 hours prior to RITUXAN infusion.

557

558 First Infusion: The RITUXAN solution for infusion should be administered
559 intravenously at an initial rate of 50 mg/hr. RITUXAN should not be mixed
560 or diluted with other drugs. If hypersensitivity or infusion reactions do not
561 occur, escalate the infusion rate in 50 mg/hr increments every 30 minutes,
562 to a maximum of 400 mg/hr. If a hypersensitivity (non-IgE-mediated) or
563 an infusion reaction develops, the infusion should be temporarily slowed
564 or interrupted (see BOXED WARNINGS and WARNINGS). The infusion
565 can continue at one-half the previous rate upon improvement of patient
566 symptoms.

567

568 Subsequent Infusions: If the patient tolerated the first infusion well,
569 subsequent RITUXAN infusions can be administered at an initial rate of
570 100 mg/hr, and increased by 100 mg/hr increments at 30-minute intervals,
571 to a maximum of 400 mg/hr as tolerated. If the patient did not tolerate the
572 first infusion well, follow the guidelines under First Infusion.

573

574 **Stability and Storage:** RITUXAN vials are stable at 2–8°C (36–46°F). Do
575 not use beyond expiration date stamped on carton. RITUXAN vials
576 should be protected from direct sunlight. Refer to the "Preparation and
577 Administration" section for information on the stability and storage of
578 solutions of RITUXAN diluted for infusion.

579

580 HOW SUPPLIED

581 RITUXAN is supplied as 100 mg and 500 mg of sterile, preservative-free,
582 single-use vials.

Single unit 100 mg carton: Contains one 10 mL vial of RITUXAN
(10 mg/mL).

585 NDC 50242-051-21

586 Single unit 500 mg carton: Contains one 50 mL vial of RITUXAN
587 (10 mg/mL).

588 NDC 50242-053-0

590 REFERENCES

- 591 1. Valentine MA, Meier KE, Rossie S, et al. Phosphorylation of the
592 CD20 phosphoprotein in resting B lymphocytes. *J Biol Chem* 1989
593 264(19): 11282–11287.
- 594
- 595 2. Einfeld DA, Brown JP, Valentine MA, et al. Molecular cloning of the
596 human B cell CD20 receptor predicts a hydrophobic protein with
597 multiple transmembrane domains. *EMBO J* 1988 7(3):711–717.
- 598
- 599 3. Anderson KC, Bates MP, Slaughenhaupt BL, et al. Expression of
600 human B cell-associated antigens on leukemias and lymphomas: A
601 model of human B cell differentiation. *Blood* 1984 63(6):1424–1433.
- 602

- 603 4. Tedder TF, Boyd AW, Freedman AS, et al. The B cell surface
604 molecule B1 is functionally linked with B cell activation and
605 differentiation. *J Immunol* 1985 135(2):973–979.
606
- 607 5. Tedder TF, Zhou LJ, Bell PD, et al. The CD20 surface molecule of B
608 lymphocytes functions as a calcium channel. *J Cell Biochem* 1990
609 14D:195.
610
- 611 6. Press OW, Applebaum F, Ledbetter JA, Martin PJ, Zarling J, Kidd P,
612 et al. Monoclonal antibody 1F5 (anti-CD20) serotherapy of human
613 B-cell lymphomas. *Blood* 1987 69(2):584–591.
614
- 615 7. Reff ME, Carner C, Chambers KS, Chinn PC, Leonard JE, Raab R, et
616 al. Depletion of B cells in vivo by a chimeric mouse human
617 monoclonal antibody to CD20. *Blood* 1994 83(2):435–445.
618
- 619 8. Demidem A, Lam T, Alas S, Hariharan K, Hanna N, and Bonavida B.
620 Chimeric anti-CD20 (IDEC-C2B8) monoclonal antibody sensitizes a B
621 cell lymphoma cell line to cell killing by cytotoxic drugs. *Cancer*
622 *Biotherapy & Radiopharmaceuticals* 1997 12(3):177–186.
623
- 624 9. Maloney DG, Liles TM, Czerwinski C, Waldichuk J, Rosenberg J,
625 Grillo-López A, et al. Phase I clinical trial using escalating single-dose

626 infusion of chimeric anti-CD20 monoclonal antibody (IDEC-C2B8) in
627 patients with recurrent B-cell lymphoma. *Blood* 1994 84(8):2457–
628 2466.

629

630 10. Berinstein NL, Grillo-López AJ, White CA, Bence-Bruckler I, Maloney
631 D, Czuczman M, et al. Association of serum Rituximab (IDEC-C2B8)
632 concentration and anti-tumor response in the treatment of recurrent
633 low-grade or follicular non-Hodgkin's lymphoma. *Annals of Oncology*
634 1998, 9:995–1001.

635

636 11. Maloney DG, Grillo-López AJ, Bodkin D, White CA, Liles T-M,
637 Royston I, et al. IDEC-C2B8: Results of a phase I multiple-dose trial
638 in patients with relapsed non-Hodgkin's lymphoma. *J Clin Oncol* 1997
639 15(10):3266–3274.

640

641 12. Maloney DG, Grillo-López AJ, White CA, Bodkin D, Schilder RJ,
642 Neidhart JA, et al. IDEC-C2B8 (Rituximab) anti-CD20 monoclonal
643 antibody therapy in patients with relapsed low-grade non-Hodgkin's
644 lymphoma. *Blood* 1997 90(6):2188–2195.

645

646 13. McLaughlin P, Grillo-López AJ, Link BK, Levy R, Czuczman MS,
647 Williams ME, et al. Rituximab chimeric anti-CD20 monoclonal
648 antibody therapy for relapsed indolent lymphoma: half of patients

649 respond to a four-dose treatment program. *J Clin Oncol* 1998
650 16(8):2825–2833.

651

652 14. Piro LD, White CA, Grillo-López AJ, Janakiraman N, Saven A, Beck
653 TM, et al. Extended Rituximab (anti-CD20 monoclonal antibody)
654 therapy for relapsed or refractory low-grade or follicular non-Hodgkin's
655 lymphoma. *Annals of Oncology* 1999 10:655–661.

656

657 15. Davis TA, Grillo-López AJ, White CA, McLaughlin P, Czuczman MS,
658 Link BK, Maloney DG, Weaver RL, Rosenberg J, Levy R. Rituximab
659 anti-CD20 monoclonal antibody therapy in non-hodgkin's lymphoma:
660 safety and efficacy of re-treatment. *J Clin Oncol* 2000 18(17):
661 3135–3143.

662

663 16. Anhalt GJ, Kim SC, Stanley JR, Korman NJ, Jabs DA, Kory M, Izumi
664 H, Rattie H, Mutasim D, Ariss-Abdo L, Labib RS. Paraneoplastic
665 Pemphigus, an autoimmune mucocutaneous disease associated with
666 neoplasia. *NEJM* 1990 323(25): 1729–1735.

667

668 17. National Institutes of Health (US), National Cancer Institute. Common
669 Toxicity Criteria. [Bethesda, MD.] : National Institutes of Health,
670 National Cancer Institute; c1998. 73p.

671

Jointly Marketed by:	
IDEC Pharmaceuticals Corporation	Genentech, Inc.
11011 Torreyana Road	1 DNA Way
San Diego, CA 92121	South San Francisco, CA 94080-4990
© 2001 IDEC Pharmaceuticals Corporation and Genentech, Inc.	

672

Revised December 2001